

Two-photon decay of pseudoscalar mesons in hot hadronic matter

T. Hashimoto

Department of Applied Mathematics, Faculty of Engineering Science, Osaka University, Toyonaka, Osaka 560, Japan

K. Hirose

Teikoku Womens University, Moriguchi, Osaka 570, Japan

T. Kanki

Faculty of Science and Technology, Kinki University, Higashi-Osaka, Osaka 577, Japan

O. Miyamura

Department of Applied Mathematics, Faculty of Engineering Science, Osaka University, Toyonaka, Osaka 560, Japan

(Received 11 August 1987)

Possible enhancement of coupling of π^0 to two photons at finite temperature is discussed in connection with second-order or weakly first-order chiral phase transition.

In this Brief Report we discuss the possibility that two-photon decay of pseudoscalar mesons at temperatures close to chiral restoration are enhanced if transition is second or weak first order.

It is well known that the PCAC (partial conservation of axial-vector current) relation with an anomaly,¹

$$\partial_\mu A^\mu = m_\pi^2 f_\pi \Phi_\pi - (\alpha/4\pi) F_{\mu\nu} \tilde{F}^{\mu\nu}, \quad (1)$$

leads to the formula of $\pi^0 \rightarrow 2\gamma$ width as²

$$\Gamma(\pi^0 \rightarrow 2\gamma) = (64\pi^3)^{-1} (\alpha/f_\pi)^2 m_\pi^3. \quad (2)$$

As the temperature independence of the anomaly has been proved,³ the second term of Eq. (1) is not modified at finite temperature. On the other hand, the coupling to axial-vector current, f_π , would receive finite-temperature correction. If the chiral transition is second order, the pion decay constant f_π is expected to vanish at the tem-

perature of chiral restoration T_χ . Therefore formula (2) suggests the enhancement of $\pi^0 \rightarrow 2\gamma$ decay at temperature close to T_χ if the pion mass remains finite.

In order to examine the above heuristic observation we study this process by an effective model which reveals chiral restoration at finite temperature. We also examine the effect of finite current-quark mass in the model.

First we calculate the triangle diagram as depicted in Fig. 1 by using the Matsubara Green's function. Here the internal line means the constituent quark with mass M and meson-constituent-quark coupling is γ_5 type (coupling constant G_{CQ}). The pion is assumed to be fixed to the rest system of thermal medium. Then we have

$$F = \epsilon_{\mu\nu\rho\sigma} e_\mu(k_1) e_\nu(k_2) k_{1\rho} k_{2\sigma} 4e^2 H, \quad (3)$$

where

$$H = G_{CQ} M T \sum_n \int \frac{d\mathbf{p}}{(2\pi)^3} \{(\mathbf{p}^2 + E_n^2 + M^2)[(\mathbf{p} - \mathbf{k})^2 + (E_n - k)^2 + M^2][\mathbf{p}^2 + (E_n - 2k)^2 + M^2]\}^{-1}, \quad (4)$$

here $E_n = 2\pi T(n + \frac{1}{2})$ and $k = |\mathbf{k}| = m_\pi/2$.

These expressions lead to 2γ decay width as

$$\Gamma_{2\gamma} = (64\pi^3)^{-1} m_\pi^3 (\alpha G_{CQ}/M)^2 I(m_\pi, M, T)^2, \quad (5)$$

where $I(m_\pi, M, T)$ is given by

$$I(m_\pi, M, T) = \int_M^\infty dE (2M^2/E)(E^2 - m_\pi^2/4)^{-1} \times \ln[(E + p)/M] \tanh(E/T). \quad (6)$$

The factor $I(m_\pi, M, T)$ represents the explicit temperature dependence of the triangle diagram at finite temperature. If masses m_π and M are fixed, I is a decreasing function of temperature and reveals thermal suppression due to interference with induced decay by the medium.

However, in Eqs. (4)–(6), we consider that masses m_π , M , and the coupling constant G_{CQ} are temperature dependent.

If the constituent-quark mass decreases near T_χ , the quark loop is enhanced as seen by the factor $(\alpha G_{CQ}/M)^2$ in Eq. (5).

In fact, if the Goldberger-Treiman relation for the constituent quark holds at finite temperature, i.e.,

$$g_A m = G_{CQ} f_\pi, \quad (7)$$

Eq. (5) is expressed as

$$\Gamma_{2\gamma} = (64\pi^3)^{-1} m_\pi^3 (\alpha g_A / f_\pi)^2 I(m_\pi, M, T)^2, \quad (8)$$

where g_A is the axial-vector coupling constant of the con-




FIG. 1. A triangle diagram for $\pi^0 \rightarrow 2\gamma$ decay at finite temperature. Internal lines are the Matsubara Green's functions for the constituent quark.

stituent quark. As g_A has a value around unity, vanishing of f_π in the case of second-order chiral restoration seems to enhance $\Gamma_{2\gamma}$. However, the validity of the Goldberger-Treiman relation at finite temperature is not obvious and should be examined. In a successful model of the Nambu–Jona-Lasinio type studied by Hatsuda and Kunihiro,⁴ the relation holds in the limit of vanishing current-quark mass. But in the case of finite current-quark mass, the relation is violated and in fact G_{CQ} also vanishes at T_χ . Based on the latter fact, decoupling of the pion from matter near critical temperature has been stressed.⁵ Thus a systematic calculation of temperature dependence of relevant masses and couplings to evaluate Eq. (5) is required.

Here we perform a calculation following their model.⁵ In the model, the Lagrangian is given by

$$L = \bar{q}(i\gamma^\mu \partial_\mu - M_0)q + g[(\bar{q}q)^2 + (\bar{q}i\gamma_5\tau q)^2], \quad (9)$$

where M_0 is the current-quark mass. The constituent-quark mass is determined from a gap equation as

$$1 - M_0/M = (3g/\pi^3)T \sum \int_{|p| < \Lambda} d\mathbf{p} \{ \mathbf{p}^2 + [(2n+1)\pi T]^2 + M^2 \}^{-1}. \quad (10)$$

Values of coupling constant g and cutoff Λ are adjusted to reproduce low-energy data such as f_π and m_π at $T=0$. The pion mass m_π and pion–constituent-quark coupling G_{CQ} are obtained by pole and residue of chain approximation of quark–antiquark scattering amplitude in a pseudoscalar channel of the form

$$\Pi_0^R(\omega, \mathbf{p}, T) / [1 - 2g\Pi_0^R(\omega, \mathbf{p}, T)], \quad (11)$$

where Π_0^R is the lowest-order contribution of the retarded




FIG. 2. A diagram for Π_0^R (Ref. 4). Lines correspond to the constituent quarks.

correlator for $\bar{q}(i\gamma_5\tau)q$ shown in Fig. 2. After determining m_π , M , and G_{CQ} by these steps we can calculate Eq. (5).

Results of the temperature dependence of relevant parameters are shown in Fig. 3(a). We see an increase of f_π^{-1} and G_{CQ}/M and an approximate validity of the Goldberger-Treiman relation except for the region very close to the vanishing point of f_π where $m_\pi = 2M$. Following the authors of Ref. 5 we define T_χ by this point. The coupling-mass ratio G_{CQ}/M quickly turns over when the temperature comes very close to T_χ . This is due to a finite current-quark mass which leads to vanishing of G_{CQ} at the same point.

In Fig. 3(b), the resultant two-photon width by Eq. (5)




FIG. 3. (a) Pion mass m_π , constituent-quark mass M , coupling-to-mass ratio G_{CQ}/M , and inverse of pion decay constant f_π^{-1} at finite temperature in the model of Refs. 4 and 5. Values of four-fermion coupling constant g , bare quark mass, and cutoff are 0.214 fm^2 , 5.5 MeV , and 631 MeV , respectively. (b) Two-photon decay width at finite temperature. $\Gamma_{2\gamma}$ [Eq. (8)] is also presented for comparison.

is presented. We also show the temperature dependence of $I(m_\pi, M, T)^2$ to see the effect of thermal suppression and the two-photon decay width by Eq. (8) for a case in which the Goldberger-Treiman relation holds (g_A is fixed to be unity).

The width grows up near T_χ in spite of the decreasing behavior of $I(m_\pi, M, T)^2$. In the chiral limit this diverges at the critical point as indicated by the calculation of Eq. (8). In the case of finite current-quark mass, it reaches a maximum. [It is noted that a zero of G_{CQ} is canceled by a singularity of $I(m_\pi, M, T)$ at T_χ .] Thus we observe the enhancement of coupling of π^0 to photons near T_χ in contrast with decoupling from matter in this model.

The following conclusion is obtained for the enhancement of the two-photon decay of π^0 near critical temperature of chiral restoration. A decrease in the mass of the constituent quark leads to an enhancement of the quark loop effect. This feature is expected if chiral restoration is a second-order or weakly first-order transition. In an example of the calculation of the Nambu–Jona-Lasinio–type model, the enhancement overwhelms thermal suppression and the decrease of the pion–constituent-quark coupling. On this point, it is remarked that the model does not simulate a deconfining transition. However, lattice Monte Carlo results indicate

the simultaneous occurrence of a chiral and deconfining transition.⁶ Thus decoupling of the pion from the quark might be affected by dynamical singularities of liberated quarks. In this regard, it is interesting that recent Monte Carlo results for shielding the mass at finite temperature has shown a clear pionic pole in the neighborhood of the transition point.⁷

Finally we mention phenomenological implications. It is an interesting challenge to observe hadron spectroscopy near the critical temperature in a high-energy collision experiment. In order to extract information of hot hadrons in a fireball, use of a penetrating probe such as the lepton or photon is appropriate.^{8,9} The photon pair is especially adequate to observe the mass shift of η' and η which is expected from effective U_1 recovery at high temperature.¹⁰ A problem in this case might be that the signal is very weak because of the shortness of the hot era of the hadronic fireball in an actual collision.¹¹ However the enhancement of two-photon decay near critical temperature discussed in this paper intensifies the signal when it also occurs for η' and η .

We would like to thank Dr. T. Hatsuda and Dr. T. Kunihiro for useful discussions.

¹S. Adler, Phys. Rev. **177**, 2426 (1969); J. S. Bell and R. Jackiw, Nuovo Cimento **60**, 47 (1969).

²R. Jackiw, *Lectures on Current Algebra and Its Application* (Princeton University Press, Princeton, NJ, 1972).

³H. Itoyama and A. H. Mueller, Nucl. Phys. **B218**, 349 (1983).

⁴T. Hatsuda and T. Kunihiro, Phys. Lett. **145B**, 7 (1984).

⁵T. Hatsuda and T. Kunihiro, Phys. Lett. **B 185**, 304 (1987); see, also, V. Bernard, Ulf-G. Meissner, and I. Zahed, MIT Report No. CTP 1446, 1987 (unpublished).

⁶Simulation for QCD with two light-quark flavors: M. Fukugi-

ta, S. Ohta, Y. Oyanagi, and A. Ukawa, Phys. Rev. Lett. **58**, 2515 (1987).

⁷C. DeTar and J. B. Kogut, Phys. Rev. Lett. **59**, 399 (1987).

⁸T. Hashimoto, O. Miyamura, K. Hirose, and T. Kanki, Phys. Rev. Lett. **57**, 2123 (1986).

⁹T. Matsui and H. Satz, Phys. Lett. **B 178**, 416 (1986).

¹⁰R. D. Pisarski and F. Wilczek, Phys. Rev. **D 29**, 338 (1984).

¹¹T. Hashimoto, K. Hirose, T. Kanki and O. Miyamura, Nucl. Phys. A (to be published).