

SECTION 8 OR SOMETHING

19 OCT 2012

ANNOUNCEMENTS:

- MON OH HELD BY JHVAM

~~Q&A~~ SOME REMARKS ABOUT HW 7

HW 4.2 $B(t) = B_0 \hat{z}$ "ASSUME AXIAL SYM"

CALL A "ASSUMING $\phi = 0$ "

↳ this is implicitly a gauge choice!

AXIAL SYM \rightarrow WANT YOU TO PICK CIRCULAR COEFFICIENT PATH s.t. $A \sim \hat{\phi}$. CHOICE OF A, COULD HAVE HAD UN COMP IN $\hat{\phi}$, $\hat{\phi}$ DIR. BUT, $\hat{\phi}$ COMP NOT COMPATIBLE w/ $\phi = 0$.

IN GENERAL (for $A \sim A_\phi \hat{\phi} + A_\psi \hat{\psi}$), WOULD NEED TO USE FARADAY'S LAW $\nabla \cdot E = -\nabla^2 \phi - \frac{1}{c} \dot{A}$ TO DETERMINE ϕ .

BUT THEN THE QUESTION IS STUPID B/C IT ASKS TO FIND ~~THE~~ E $\nabla \cdot E$ THEN CONFIRM FARADAY.

BUT YOU NEED FARADAY TO (IN GENERAL GAUGE) FIND E.

NEW ("macroscopic") MAXWELL EQNS

$$\nabla \cdot \underline{D} = 4\pi \rho_f$$

$$\nabla \cdot \underline{B} = 0$$

$$\nabla \times \underline{E} = -\dot{\underline{B}}$$

$$\nabla \times \underline{H} = \dot{\underline{D}} + \underline{J}_f$$

$$\underline{D} = \underline{E} + 4\pi \underline{P} = \epsilon \underline{E}$$

$$\underline{H} = \underline{B} - 4\pi \underline{M} = \frac{1}{\mu} \underline{B}$$

note: $\nabla \cdot \underline{H} = -4\pi \nabla \cdot \underline{M}$
not "analogous to \underline{B} "


DIVERGENCE LAWS \rightarrow (DIS) CONTINUITY of NORMAL COMP.

CURL LAWS \rightarrow (DIS) CONTINUITY of TANGENTIAL

eg. in absence of FREE CHARGE,

D_{\perp} is continuous $\Rightarrow E = \frac{1}{\epsilon} D$ is NOT.

HPM 1.13



WHAT IS $E(0)$?

INTEGRATE COULOMB FIELD of POINT SURFACE CHARGE σ_b . (or P_b)

$$P_b = \underline{n} \cdot \underline{P} = P \cos \theta$$

$$E(\underline{r}) = \int \frac{P(s)}{|\underline{r}-s|^2} \hat{(\underline{r}-s)} d^3s \rightarrow \int \frac{P_b}{a^2} \hat{(\underline{r})} (2\pi) a^2 d(\cos \theta)$$

GO AHEAD + PROJECT ON E_z , ONLY NONZERO COMP

$$E_z(0) = \int \frac{1}{a^2} P \cos \theta \cdot \underbrace{\hat{(\underline{r})} \cdot \hat{(\underline{z})}}_{-\cos \theta} (2\pi) a^2 d(\cos \theta)$$


$$E_z(0) = -2\pi P \int_{-1}^1 \cos^2 \theta d(\cos \theta) = \boxed{-\frac{4\pi}{3} P}$$

$$\underbrace{\int_{-1}^1 \cos^2 \theta d(\cos \theta)}_{\frac{1}{3} u^3 \Big|_{-1}^1 = \frac{2}{3}}$$

$$\Rightarrow \boxed{E(0) = -\frac{4\pi}{3} P}$$


b) DOES THIS LOOK REASONABLE? WHAT ABOUT OTHER $r < a$?

CLAIM: COMPARE TO OVERLAP OF OPP CHARGED UNIFORM SPHERES



CHOOSE $P \neq q$ s.t. OVERLAP CONFIG MATCHES PART (a)

NOTE: PART 2 HAS TOTAL DIPOLE MOMENT $\boxed{P \frac{4\pi a^3}{3}} = P_{TOT}$



~ •

s.t. SUPERIMPOSED CONFIG HAS DIPOLE MOMENT

$$P = q \delta \quad \text{w/ } q = \frac{4\pi a^3}{3} \rho$$

THE DIPOLE MOMENTS MATCH WHEN $P = p \delta$

CLAIM: IN A UNIFORM SPHERE $E(r) = \frac{4\pi}{3} P r$

check!

then: ~~the~~ $E = \frac{4\pi}{3} P \left[(r - \frac{R}{2}) - (r + \frac{R}{2}) \right]$
 $= -\frac{4\pi}{3} P R$

$$\boxed{E = -\frac{4\pi}{3} P R}$$

what we found @ $r=0$

in fact, E is constant everywhere inside sphere!

ALTERNATE DERIVATION: SPHERICAL HARMONICS

$$P_z = P \cdot P_1(\cos \theta)$$

(c) NOW CONSIDER A POLARIZED MEDIUM w/ A SPHERICAL CAVITY CARVED OUT. WHAT IS E_c IN THE CAVITY w/RT E IN THE MEDIUM? (E IS EXTERNAL)


SAME TRICK: SUPERIMPOSE A POLARIZED SPHERE

w/ OP POL TO DIELECTRIC. ie POLARIZE $P_{\text{sup}} = -P$

$$\underline{E}_{\text{ext}} + \underline{E}_{\text{sup}} = \underline{E} - \frac{4\pi}{3} P_{\text{sup}} = \boxed{\underline{E} + \frac{4\pi}{3} P}$$

note: we are ignoring dipole contribution!

EXAMPLES: MIDTERM w/ DIELECTRIC



$$\begin{aligned} & \cancel{= \sum C_l r^l P_l} \\ & = (C_l r^l + D_l \frac{1}{r^{l+1}}) P_l \\ & = A_l r^l P_l \end{aligned}$$

BC: CONTINUITY OF $E_{||}$: \oint IS CONTINUOUS

DISCONTINUITY OF D_{\perp} : $\epsilon' \frac{\partial \phi'}{\partial r} - \epsilon \frac{\partial \phi}{\partial r} = 4\pi\sigma$

START @ BOUNDARY $r=b$, SIMPLIFY.

$$\begin{aligned} \phi(b) = 0 & \Rightarrow \boxed{D_l = -C_l b^{2l+1}} \\ \phi & = \sum C_l (r^l - \frac{b^{2l+1}}{r^{l+1}}) P_l \end{aligned}$$

NEXT SIMPLIFY: CONTINUITY @ $r=a$

$$\phi'(a) = \phi(a) \Rightarrow A_l a^l = C_l (a^l - \frac{b^{2l+1}}{a^{l+1}})$$

$$\boxed{A_l = C_l \left(1 - \left(\frac{b}{a}\right)^{2l+1}\right)}$$

so far: SANS!

HARD BC

$$\text{at } r=a, \quad \underline{\underline{\epsilon \partial_r \phi(a) - \partial_r \phi'(a) = -4\pi\sigma_0}}$$

$$\begin{aligned} & \sum_l \epsilon C_l \left[l a^{l-1} + (l+1) \frac{b^{2l+1}}{a^{l+2}} \right] P_l - \sum_l C_l \left[l a^{l-1} + (l+1) \frac{b^{2l+1}}{a^{l+2}} \right] P_l \\ & = -4\pi\sigma_0 \left[\frac{1}{3} P_0 + P_1 + \frac{2}{3} P_2 \right] \end{aligned}$$

like: $\underline{\underline{\epsilon C_l l a^{l-1} + \epsilon C_l (l+1) \frac{b^{2l+1}}{a^{l+2}}} - C_l l a^{l-1} + C_l l \frac{b^{2l+1}}{a^{l+2}} = \dots}$

$$C_l \left[(\epsilon+1)l + 1 \right] \frac{b^{2l+1}}{a^{l+2}} + (\epsilon-1) C_l l a^{l-1} = \dots$$

SO MATCHING COMPONENTS:

$$C_0 \frac{b^0}{a^2} = -4\pi\sigma_0 \frac{1}{3}$$

\rightarrow

$$C_0 = (-4\pi\sigma_0) \frac{1}{3} \frac{1}{b^0/a^2}$$

$$C_1 \left[(\epsilon+1)2 + 1 \right] \frac{b^3}{a^3} + (\epsilon-1) C_1 \cdot 1 = -4\pi\sigma_0$$

$$\hookrightarrow C_1 \left[(\epsilon+2) \frac{b^3}{a^3} + (\epsilon-1) \right] = -4\pi\sigma_0$$

$$C_1 = -4\pi\sigma_0 \left[(\epsilon+2) \frac{b^3}{a^3} + (\epsilon-1) \right]^{-1}$$

$$C_2 \left[(\epsilon+1)3 + 1 \right] \frac{b^5}{a^4} + (\epsilon-1) C_2 2a = -4\pi\sigma_0 \frac{2}{3}$$

$$C_2 = -4\pi\sigma_0 \frac{2}{3} \left[(\epsilon+3) \frac{b^5}{a^4} + 2(\epsilon-1)a \right]^{-1}$$

INDUCED ~~CHARGE~~ CHARGE @ $r=b$

$$-\partial_r \phi(b) = -4\pi\sigma^{\text{INDUCED}} \rightarrow \sigma = \frac{\partial_r \phi(b)}{4\pi}$$

$$\partial_r \phi(r) = \sum c_\ell \left[\ell r^{\ell-1} + (\ell+1) \frac{b^{2\ell+1}}{r^{\ell+2}} \right] P_\ell$$

$$\frac{\partial_r \phi(b)}{4\pi} = -\sigma_0 \left[\frac{4}{3} \frac{a^2}{b} \left(0 + \frac{1}{b} \right) P_0 \right.$$

$$+ \left[(\epsilon+2) \frac{b^3}{a^3} + (\epsilon-1) \right]^{-1} (1+2) P_1$$

$$\left. + \frac{2}{3} \left[(\epsilon+3) \frac{b^5}{a^4} + 2(\epsilon-1)a \right]^{-1} P_2 \right]$$

UUGLY! but can see: ϵ affects different multipoles differently.