

Distinguishing s-channel resonances At the ILC



Steve Godfrey & Alex Tomkins
Carleton University

ILC Workshop, Snowmass Colorado, August 25, 2005



- Little Higgs W_H^\pm Z_H B_H
- Extra dimensions (ADD, RS, UED...): KK excitations
 - ADD: Graviton tower exchange effective operators: $i \frac{4\lambda}{M_H^4} T^{\mu\nu} T_{\mu\nu}$
 - Randall-Sundrum Gravitons: Discrete KK graviton spectrum
- Extended gauge sectors
 - Extra U(1) factors: $E_6 \rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi$
 - Left-Right symmetric model: $SU(2)_L \times SU(2)_R \times U(1)$
- Topcolour

Many, many models

What do these models have in common?

- Almost all of these models have new s -channel structure at \sim TeV scale
- Either from extended gauge bosons or new resonances

How do we distinguish the models?



How do we distinguish them?

Start by assuming the LHC discovers single rather heavy resonance


What is it?

Tools are:


- Cross sections & Widths
- Angular Distributions
- Couplings (decays, polarization...)

On resonance production of (RS) Gravitons

Use angular distributions to test against different spin hypothesis



Measure BR's to test for Universal couplings





Davoudiasl, Hewett and Rizzo,
Phys. Rev. D63, 075004 (2001)
[hep-ph/0006041].



Interference of exchange of virtual graviton KK states with SM amplitudes

ADD:
$$i \frac{4\lambda}{M_H^4} T^{\mu\nu} T_{\mu\nu}$$

Leads to deviations in $e^+e^- \rightarrow f\bar{f}$ dependent on both λ and s/M_H



$\sqrt{s} = 5 \text{ TeV} \quad L = 1 \text{ ab}^{-1} \quad M_s = 15 \text{ TeV} \quad \lambda = \pm 1$

Can use multipole moments to distinguish spin 2 from spin 1

Rizzo: hep-ph/0208027





ID ADD Graviton Exchange

Pankov & Paver hep-ph/0501170

Suitable observables can divide possible models into subclasses

- To identify graviton exchange
- Forward-Backward Centre-Edge

asymmetries: $\sigma_{CE,FB} = \sigma_{C,FB} - \sigma_{E,FB}$




LC: 5σ **ident. reach** on $M_H = 3.5 - 5.8 \text{ TeV}$ at $\sqrt{s} = 0.5 - 1 \text{ TeV}$ and $\mathcal{L}_{\text{int}} = 500 \text{ fb}^{-1}$



Numerous difermion observables

18 di-fermion observables:

- σ^μ
- A_{FB}^μ
- A_{LR}^μ
- $A_{FB}^\mu(pol)$
- σ^τ
- A_{FB}^τ
- A_{LR}^τ
- P_τ
- R^{had}
- A_{LR}^{had}
- σ^b
- A_{FB}^b
- A_{LR}^b
- $A_{FB}^b(pol)$
- σ^c
- A_{FB}^c
- A_{LR}^c
- $A_{FB}^c(pol)$



Z' couplings

Extraction of Z' couplings
assuming $M_{Z'}$ is known from LHC

$$\sigma_{P_e^- P_e^+}^\mu$$


$$A_{FB}^\mu$$

$$A_{LR}^\mu$$

95% C.L. bounds

$L=1 \text{ ab}^{-1}$, $\Delta L=0.2\%$, $P_- = 0.8$, $P_+ = 0.6$, $\Delta P=0.5\%$

Note sign ambiguity



S. Riemann: TESLA TDR & LHC/LC Study





1. Can we resolve the sign ambiguity?

$$\sqrt{s} = 500 \text{ GeV}$$

$$\mathcal{L}_{int} = 1 \text{ ab}^{-1}$$

$$M_{Z'} = 1, 1.5, 2 \text{ TeV}$$





2. How does it change with more observables?

$$\sqrt{s} = 500 \text{ GeV} \quad \mathcal{L}_{int} = 1 \text{ ab}^{-1}$$



$$\sigma^\mu \quad A_{FB}^\mu \quad A_{LR}^\mu$$

$$\sigma^\mu \quad A_{FB}^\mu \quad A_{LR}^\mu \quad A_{FB}^\mu (pol) \quad \sigma^\tau \quad A_{FB}^\tau \quad A_{LR}^\tau \quad P_\tau$$



3. What happens for higher mass?


$$\sqrt{s} = 500 \text{ GeV} \quad \mathcal{L}_{int} = 1 \text{ ab}^{-1}$$



4. What happens with higher energy?


$$M_{Z'} = 2.5 \text{ TeV}$$

$$\sqrt{s} = 500, 800, 1000, 1500 \text{ GeV}$$




The "Peskin Plot"

$M_{Z'} = 1, 1.5, 2, 2.5$ TeV



$M_{Z'} = 1, 2, 3, 4$ TeV



$\sqrt{s} = 500$ GeV $\mathcal{L}_{int} = 1$ ab $^{-1}$



The ILC will be an extremely powerful tool for understanding a resonance discovered at the LHC